

Introduction

Vegetation remnants in the eastern Darling Downs are highly fragmented and greatly reduced in size. Since 1975 there has been up to a 60% reduction in native vegetation in the region. Little is known about habitat quality and condition of remnant and re-growth vegetation, including endangered ecosystems such as bluegrass grasslands & semi-evergreen vine thickets.

The objective of this research was to compare plant species richness, condition and habitat complexity in remnant vegetation in the study area.

Figure 1: Map of north eastern Murray-Darling Basin showing location of study area



Results & Discussion

- richness = 31-83 spp./500m²; habitat complexity = 6-17; condition = 23-31.
- significant patterns in species richness, complexity and condition across vegetation types (Fig. 2).
- grasslands - low complexity, fewer species, high condition.
- Mt Coolibah & Ironbark/Mt Coolibah woodlands - low condition, intermediate richness and complexity.
- richness-complexity strongly correlated ($R_s = 0.54$ $P < 0.001$); richness-condition poorly related ($R_s = -0.20$ $P > 0.05$).


Methods

- 43 sites were sampled across 11 vegetation types in the study area in southern Queensland (Fig. 1).
- Plant species richness was determined in a single 500 m² quadrat at each site.
- Habitat complexity¹ was derived from vegetation structure (FPC of strata, cover of litter, logs etc) and other biophysical attributes (e.g. hollows, stags etc).
- A measure of vegetation condition was derived from the summation of scores for range of attributes (Table 1).
- One-way anova compared attributes across vegetation types; Spearman Rank correlations examined relationships between attributes.

Table 1: Components of Condition Index

Component	Method
Physical disturbance Grazing, Clearing, Erosion, Weeds, Ferals, Logging, Epiphytic Growth, Compaction, Canopy Death	Attributes ranked in field (0-3; 0=low, 3=high dist) - inverse rank used to determine score (ie 0=degraded, 3=healthy)
Juvenile Density [trees] (3 classes: <1, 1-3, x3m ht)	Ranked 0=0, 1=1-10, 2=10-20, 3=>20 individuals /500m ²
Ground cover Litter Cover & Bare ground Cover	Litter 0=0%; 1=1-10%; 2=10-30%; 3=>30% cover Bare ground 0=>20%; 1=10-20%; 2=<10%; 3=0%

Figure 2: Comparison of Species Richness, Habitat Complexity and Condition across vegetation types



Acknowledgements

This project is part of the Land Use Change Project by the Land Use Research Centre and supported by a USQ Project Team Research grant. We thank the many landowners in the eastern Darling Downs for allowing access to properties. Further information: landuse@usq.edu.au

While species richness alone is not a definitive attribute of vegetation, current theory would suggest that richness would be related to condition.

Developments in assessing soil condition² and habitat quality³ may prove to work well where there are suitable reference sites; however, broader application of condition indices, particularly in regions that are highly modified, may require considerable internal calibration.

Conclusions

There is a need for simple, robust and meaningful on-ground indicators of the impacts of human activities in remnants.

Methods for habitat complexity have been developed for some time and have been rigorously tested over a range of ecosystems.

However, the notion of habitat condition remains rather vague and largely untested. This is despite a growing acceptance of this attribute as a decision tool by land managers.

Clearly, further development of methods for the accurate quantification of condition is necessary.

References

1. Isaacson A.E., & Culling P.C. (1979) Habitat preferences of mammals inhabiting heathlands of temperate coastal, montane and alpine regions of southeastern Australia. In *Ecosystems of the World*, vol. 24, *Heathlands and Related Shrublands of the World*, (Ed. R.L. Specht.) Elsevier, Amsterdam.
2. Ludwig J.A. & Tonger D.J. (1997) A landscape approach to Rangeland Ecology. In *Landscape Ecology Function and Management: Principles from Australia's Rangelands*, (Eds J. Ludwig, D. Tonger, D. Freudenberger, J.C. Noble & K. Hodgkinson) CSIRO, Melbourne.
3. Parkes D., Benall G. & Chell D. (2005) Assessing the quality of native vegetation: the 'habitat condition' approach. *Ecological Management & Restoration* 4: 529-535.